## Class X Mathematics –Standard (041) Sample Question Paper 2019-20

## Max. Marks: 80

**Duration : 3 hrs** 

## General Instructions:


- (i) All the questions are compulsory.
- (ii) The question paper consists of 40 questions divided into 4 sections A, B, C, and D.
- (iii) Section A comprises of 20 questions of 1 mark each. Section B comprises of 6 questions of 2 marks each. Section C comprises of 8 questions of 3 marks each. Section D comprises of 6 questions of 4 marks each.
- (iv) There is no overall choice. However, an internal choice has been provided in two questions of 1 mark each, two questions of 2 marks each, three questions of 3 marks each, and three questions of 4 marks each. You have to attempt only one of the alternatives in all such questions.
- (v) Use of calculators is not permitted.


## SECTION A


	- Q 10 are multiple choice questions. Select the most appropriate answer from the en options.	
1	The decimal representation of $\frac{11}{2^3 \times 5}$ will a) terminate after 1 decimal place b) terminate after 2 decimal places c) terminate after 3 decimal places d) not terminate	1

2	Consider t	ne following	frequency d	istribution o	f the heights	of 60 stude	ents of a class	1
	Height (in	150-155	155-160	160-165	165-170	170-175	175-180	
	cm)							
	No of	15	13	10	8	9	5	
	students							
	The upper	limit of the r	nedian class	s in the give	n data is			
	a) 165							
	b) 155							
	c) 160 d) 170							
	d) 170	)						
3		of smallest ty	wo digit com	posite numb	per and sma	llest compos	site number is	1
	a) 12 b) 4							
	c) 20							
	d) 44							
				$\mathbf{R}$	es	ulf.	in	
						UIL		
4	For which	value(s) of n	, will the line	es represent	ted by the fo	llowing pair	of linear	1
		be parallel	,		,			
				3x - y -				
			voont 10	6x - 2y -	p = 0			
	b) 10	eal values e						
	c) 5/2							
	d) 1/2							
L								


5	If triangle ABC is right angled at C, then the value of sec (A+B) is a) 0 b) 1 c) $\frac{2}{\sqrt{3}}$	1
	d) not defined	
6	If $sin\theta + cos\theta = \sqrt{2}cos\theta$ , $(\theta \neq 90^\circ)$ then the value of $tan\theta$ is	1
	a) $\sqrt{2} - 1$ b) $\sqrt{2} + 1$	
	c) $\sqrt{2}$	
	d) $-\sqrt{2}$	
7	Civen that since $\sqrt{3}$ and $\cos\theta$ . O then the value of $\theta$ with	1
-	Given that $sin\alpha = \frac{\sqrt{3}}{2}$ and $cos\beta = 0$ , then the value of $\beta - \alpha$ is	
	a) 0° b) 90°	
	c) 60°	
	d) 30°	
8	The point which divides the line segment joining the points (8, – 9) and (2, 3) in ratio 1 : 2 internally lies in the	1
	a) I quadrant	
	b) II quadrant	
	c) III quadrant d) IV quadrant	
	u) iv quadrant	
9	The distance of the point P $(-3, -4)$ from the x-axis (in units) is	1
	a) 3	
	b) -3 c) 4	
	d) 5	
L		








25	Jayanti throws a pair of dice and records the product of the numbers appearing on the dice. Pihu throws 1 dice and records the squares the number that appears on it. Who has the better chance of getting the number 36? Justify?	2
	OR	
	<ul> <li>An integer is chosen between 70 and 100, Find the probability that it is</li> <li>(a) a prime number</li> <li>(b) divisible by 7</li> </ul>	
26	Isha is 10 years old girl. On the result day, Isha and her father Suresh were very happy as she got first position in the class. While coming back to their home, Isha asked for a treat from her father as a reward for her success. They went to a juice shop and asked for two glasses of juice.	2
	Aisha, a juice seller, was serving juice to her customers in two types of glasses. Both the glasses had inner radius 3cm. The height of both the glasses was 10cm.	
	First type: A Glass with hemispherical raised bottom.	
	ClassResult.in	
	Second type: A glass with conical raised bottom of height 1.5 cm.	
	Isha insisted to have the juice in first type of glass and her father decided to have the juice in second type of glass. Out of the two, Isha or her father Suresh, who got more quantity of juice to drink and by how much?	
	Section C	
27	Given that $\sqrt{5}$ is irrational, prove that $2\sqrt{5} - 3$ is an irrational number.	3
	OR	
	If HCF of 144 and 180 is expressed in the form 13m-16. Find the value of m.	



OREvaluate: $\frac{cos^2(45^\circ+\theta)+cos^2(45^\circ-\theta)}{\tan(60^\circ+\theta)\times\tan(30^\circ-\theta)} + (cot30^\circ+sin 90^\circ) \times (tan60^\circ-sec 0^\circ)$ 33Sides of a right triangular field are 25m, 24m and 7m. At the three corners of the field, a cow, a buffalo and a horse are tied separately with ropes of 3.5 m each to graze in the field. Find the area of the field that cannot be grazed by these animals.	32	<ul> <li>(i) Who will travel more distance, Seema or Aditya, to reach to their hometown?</li> <li>(ii) Seema and Aditya planned to meet at a location D situated at a point D represented by the mid-point of the line joining the points represented by the point D.</li> <li>(iii) Find the area of the triangle formed by joining the points represented by A, B and C.</li> </ul>	3
$\frac{\cos^{2}(45^{\circ}+\theta) + \cos^{2}(45^{\circ}-\theta)}{\tan(60^{\circ}+\theta) \times \tan(30^{\circ}-\theta)} + (\cot 30^{\circ} + \sin 90^{\circ}) \times (\tan 60^{\circ} - \sec 0^{\circ})$ 33 Sides of a right triangular field are 25m, 24m and 7m. At the three corners of the field, a cow, a buffalo and a horse are tied separately with ropes of 3.5 m each to graze in the		OR	
cow, a buffalo and a horse are tied separately with ropes of 3.5 m each to graze in the			
	33	cow, a buffalo and a horse are tied separately with ropes of 3.5 m each to graze in the	3



36			•	rallel to one other two sig		•		e other two	4
37				) km at a un nutes less f	•		•		
					OR				
	Solve the f	ollowing ec	quation:						
	$\frac{1}{x}$ -	$\frac{1}{x-2} = 3, x$	;≠0,2						
38	lower and	upper ends	s as 20 m a	a frustum of and 50 m re ate of Rs. 70	spectively.	Find the co	ost of petro	l which can	
	OR								
		ond which i		f 15km/hou g and 44m w	r through a				
39	cuboidal po pond rise b The angle 30 second	ond which i by 21cm? of elevatio s, the angl	s 50m long <b>as</b> n of an air e of eleva	f 15km/hou	r through a vide. In what e point on the es 30°. If the	time will the ground	is 60°. Afte	water in the	4
	cuboidal po pond rise b The angle 30 second height of 30	ond which i by 21cm? of elevatio s, the angl $000\sqrt{3}$ m, f	s 50m long as n of an air e of eleva ind the spe	f 15km/hou g and 44m w SR plane from a tion become eed of the air	r through a ride. In what e point on the rplane.	time will the ground e airplane	is 60 <sup>0</sup> . After	water in the	4
39 40	cuboidal po pond rise b The angle 30 second height of 30	ond which i by 21cm? of elevatio s, the angl $000\sqrt{3}$ m, f	s 50m long as n of an air e of eleva ind the spe	f 15km/hou g and 44m w SR plane from a tion become	r through a ride. In what e point on the rplane.	time will the ground e airplane	is 60 <sup>0</sup> . After	water in the	4